



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Effects of antibiotics on -toxin levels during *Staphylococcus aureus* culture: implications for the protection of chondrocytes in a model of septic arthritis.

Citation for published version:

Miller, RP, Berlouis, ME, Hall, AG, Simpson, AHRW, Smith, IDM & Hall, A 2019, 'Effects of antibiotics on -toxin levels during *Staphylococcus aureus* culture: implications for the protection of chondrocytes in a model of septic arthritis.', *Cartilage*. <https://doi.org/10.1177/1947603519828433>

Digital Object Identifier (DOI):

[10.1177/1947603519828433](https://doi.org/10.1177/1947603519828433)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Cartilage

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Effects of antibiotics on α -toxin levels during *Staphylococcus aureus* culture: implications for the protection of chondrocytes in a model of septic arthritis.

R. P. Miller, M. E. Berlouis, A. G. Hall, A. H. R.W. Simpson¹,
I. D. M. Smith² and A. C. Hall*.

Centre for Integrative Physiology, Deanery of Biomedical Sciences, University of Edinburgh, Edinburgh
EH8 9AG, Scotland, U.K.

¹Musculoskeletal Research Unit, Department of Orthopaedic Surgery, University of Edinburgh, Chancellor's
Building, 49 Little France Crescent, Edinburgh EH16 4SB, U.K.

²Institute of Infection, Immunity and Inflammation, University of Glasgow, Glasgow, G12 8TA, Scotland,
U.K.

*Author for correspondence:

Andrew C. Hall,
Centre for Integrative Physiology,
Deanery of Biomedical Sciences,
University of Edinburgh,
Hugh Robson Building,
George Square,
Edinburgh, EH8 9AG,
Scotland, U.K.

Phone: +44 (0)131 650 3263

FAX: +44 (0)131 650 2872

e.mail: a.hall@ed.ac.uk

Running title: Antibiotics on *S. aureus* α -toxin levels

Abstract

Objective: Septic arthritis results from joint infection by *Staphylococcus aureus* which produces potent alpha (α)-toxin causing cell death, potentially leading to permanent cartilage damage. Treatment is by joint irrigation and antibiotics, although it is unclear if, following treatment with antibiotics which cause bacterial lysis, there is release of additional stored (α)-toxin.

Design: A rabbit erythrocyte haemolysis assay was optimised to assess biologically-active α -toxin from cultured *S. aureus* (α)-toxin strain DU5946. Haemoglobin release was measured spectrophotometrically following addition of a bacteriostatic/bactericidal antibiotic (linezolid) or a bacteriolytic antibiotic (penicillin). A bovine cartilage model of septic arthritis was utilised to test the protective effects of antibiotics against *S. aureus* infection.

Results: During *S. aureus* culture, α -toxin levels increased rapidly but the rate of rise was quickly (within 20mins) suppressed by linezolid (25 μ g/ml). Penicillin also reduced the increase in α -toxin levels, however the time course was relatively slow compared to linezolid even at high concentrations (50,000U/ml). The efficacy of penicillin (250,000U/ml) at reducing the rise in α -toxin was approximately 8% less than that of linezolid ($P<0.05$) suggesting the presence of additional toxin. This could be due to a delayed action of penicillin, and/or release of a small pool of stored α -toxin from dying bacteria. In a bovine cartilage model however, there was no difference between the protection of *in situ* chondrocytes against *S. aureus* by penicillin or linezolid ($P>0.05$).

Conclusion: The results suggested that equally effective protection of chondrocytes against *S. aureus* septic arthritis may be obtained by the bacteriostatic/bactericidal or bacteriolytic antibiotics tested.

Key words: *Staphylococcus aureus*; bacteriostatic; bactericidal; antibiotic; α -toxin; septic arthritis.

Introduction

Septic arthritis, resulting from joint inflammation secondary to infection¹, can be highly destructive leading to cartilage damage and joint failure²⁻⁴. It affects all ages with an incidence in Western Europe of 4-10 cases/100,000 persons/yr⁵. The incidence of septic arthritis is rising globally due to various factors including an ageing population, increased use of immunosuppressive agents, musculoskeletal prosthesis and surgical procedures⁵⁻⁷. Although numerous bacterial species may induce septic arthritis, *Staphylococcus aureus* accounts for 40-60% of cases^{2,3,8} with invasive healthcare-associated methicillin-resistant *S. aureus* (MRSA) infections accounting for 18 per 100,000 persons/yr⁹. The bacteria enter the synovial joint mainly by haematogenous spread⁵, and the infection results in chondrocyte death, leading to cartilage degeneration and tissue erosion^{2,3}. Chondrocyte death is particularly important as these are the only cell type capable of maintaining cartilage resilience through their regulation of extracellular matrix metabolism¹⁰. Septic arthritis can lead to osteoarthritis (OA¹¹) which contributes to the high morbidity associated with the disease, and may also lead to fatal septicaemia⁷.

S. aureus produces an array of potential virulence factors (e.g. toxins, adhesins), which contribute to colonisation and subsequent tissue destruction. The toxins may damage host cells directly or facilitate evasion of the host immune response¹². Depending on the strain, *S. aureus* may release toxins including α , β , and γ -toxins in addition to damaging agents including Pantone-Valentine leucocidin (PVL) and super-antigens (e.g. toxic shock syndrome toxin)¹³. A bovine cartilage explant model of *S. aureus*-induced septic arthritis, however reported that the 'pore forming' alpha (α)-toxin is the key damaging agent causing rapid chondrocyte death whereas the β and γ -toxins were relatively ineffectual^{14,15}. Alpha-toxin is active against a variety of mammalian cells, but has marked potency against rabbit red blood cells (RBCs) *via* the ADAM-10 receptor which is also present on human chondrocytes¹⁶⁻¹⁸. Rabbit RBCs haemolysis can be quantified spectrophotometrically and thus is a very useful and sensitive biological assay for *S. aureus* α -toxin^{19,20}.

Treatment for septic arthritis includes intravenous antibiotics with joint lavage¹, however retrospective studies indicate that some permanent cartilage damage develops in ~50% of cases²⁻⁴, thus its rapid and effective treatment is essential. When patients present, the joint is aspirated to obtain a microbiological profile of the synovial fluid. As identification of the causative micro-organism may take ~48hrs, patients are traditionally empirically commenced on intravenous antibiotics with joint lavage. Beta (β) lactam type antibiotics such as benzyl penicillin (penicillin-G) and flucloxacillin are classed as bacteriolytic as they kill the bacteria through damage to the cell wall^{21,22}. However, this may release cellular contents including *S. aureus* α -toxin potentially causing additional chondrocyte death above that occurring when the bacteria were alive. In contrast, antibiotics such as linezolid and erythromycin are bacteriostatic at low concentrations by inhibiting bacterial growth and replication, which may then be followed by their removal by the immune system^{23,24}, whereas at higher levels they are bactericidal^{24,25}. Thus these agents can be described as bacteriostatic/bactericidal. The choice of antibiotic for *S. aureus* septic arthritis treatment is important as it could influence the amount of α -toxin in the synovial space, chondrocyte death and subsequent cartilage damage.

98 Here, we have utilised two isogenic mutants (DU5946, DU5938) from the well-characterised
99 prophage-cured derivative of NCTC8325 *S. aureus* 8325-4²⁶. NCTC8325 was originally isolated from a patient
100 with joint sepsis and its lineage remains a valuable resource for *S. aureus* research²⁷. These mutants produce
101 the range of toxins^{26,28,29} and the mutations only affect the synthesis of alpha-haemolysin (α)-toxin (Hla::Em^r),
102 beta (β)-haemolysin (Hlb:: ϕ 42E) and gamma (γ)-haemolysin (Δ hlg::Tc^r). Thus, while both strains produce
103 damaging agents, the mutant strain DU5946 only produces α -toxin and not β or γ -haemolysin, whereas the
104 DU5938 mutant does not produce α , β , or γ -toxins. Levels of biologically active α -toxin during *S. aureus*
105 culture, were assessed semi-quantitatively using the sensitive rabbit haemolysis assay^{19,20}. Using this
106 technique, we tested the hypothesis that α -toxin levels in the culture medium would be higher after the addition
107 of high concentrations of a bacteriolytic antibiotic (penicillin G) compared to a bacteriostatic/bactericidal
108 antibiotic (linezolid). The results demonstrated that there was only a small (<10%) further increase in α -toxin
109 levels from *S. aureus* cultures following penicillin treatment compared to linezolid. We then tested these two
110 antibiotics (in addition to two other antibiotics, flucloxacillin (bacteriolytic) and erythromycin
111 (bacteriostatic/bactericidal)) for their ability to protect articular chondrocytes in a bovine cartilage model of
112 septic arthritis. The results suggested that both penicillin and linezolid (as well as flucloxacillin and
113 erythromycin) protected chondrocytes with equal efficacy suggesting that there was a negligible intracellular
114 α -toxin released from *S. aureus* following antibiotic treatment.
115

Materials and Methods

(a): *Bacterial strains and reagents*. Two isogenic mutants of *S. aureus* 8325-4 (DU5946, DU5938) containing mutations affecting the synthesis of alpha-haemolysin (α)-toxin (Hla::Em^f), beta-haemolysin (Hlb:: ϕ 42E) and gamma haemolysin (Δ hlg::Tc^r; Table 1) were used. The DU5946 strain produced α -toxin whereas this was not produced by the DU5938 strain, otherwise the strains were identical. Antibiotics were from Sigma-Aldrich, Irvine, UK. Fresh, filter-sterilised stock solutions of linezolid (PZ0014), Na-flucloxacillin (F0150000) and penicillin G (P3032; 1477U/mg) were prepared in dH₂O whereas erythromycin (E6376) was dissolved in ethanol-water (6mg/ml).

(b): *Preparation of bacterial pellets, supernatant samples and antibiotics*. Bacteria were stored at -80°C in 10%v/w skimmed milk (Oxoid, Basingstoke, UK). When required, they were thawed and streaked onto tryptone soya agar (TSA; Oxoid) plates containing 2 μ g/ml tetracycline (Cat. 87128; Sigma-Aldrich). Following incubation (24hrs; 37°C), tryptone soya broth (TSB; Oxoid), containing 2 μ g/ml tetracycline, was inoculated with several single bacterial colonies from the TSA plate and incubated (24hr; 37°C) with shaking. From this TSB culture, serial dilutions were performed in saline to 10⁻⁶, in order to calculate the number of colony forming units (CFU). A CFU is defined as a unit used to estimate the number of viable bacteria capable of reproducing to form a colony of the same bacterial species – the number of CFUs is therefore a measure of the number of active bacteria in 1ml of TSB cultured for 24hr. Thereafter, 100 μ l of 10⁻⁴, 10⁻⁵ and 10⁻⁶ dilutions were spread evenly onto TSA plates and incubated (37°C; 24hr). Colonies were then counted using a colony counter (Stuart[®], Bibby Scientific, Stone, UK). Bacterial counts were performed on several cultures for each strain and a count of $\sim 1 \times 10^9$ CFU/ml (range 0.8-1.25 $\times 10^9$ CFU/ml) was routinely obtained. Cultures were then centrifuged (2,000xg; 15min) and the pellets resuspended by vortexing, washing in HBSS and centrifugation (x3). The supernatant was then removed and the pellets air-dried before being stored (-20°C) and used within one week. Based on the colony counts, a bacterial pellet contained $\sim 1 \times 10^5$ CFU and the bacteria were used in the experiments described here at $\sim 0.2 \times 10^5$ CFU in HBSS. TSA plates were prepared to assess effective doses of antibiotics against *S. aureus* whose viability was assessed by counting colonies plated on TSA in the presence/absence of antibiotics following incubation overnight at 37°C. For penicillin and linezolid, doses of 1000U/ml and 25 μ g/ml respectively were sufficient. For flucloxacillin and erythromycin, doses of 2mg/ml and 10 μ g/ml respectively were adequate and no viable bacteria were detected. For some experiments where α -toxin levels were assessed by the haemolysis method, higher doses of penicillin (to 250,000U/ml) were used. Table 2 summarises the antibiotics used with doses, together with sites of action and references. All doses used were substantially greater than the MIC (minimum inhibitory concentration) in the literature (Table 2).

(c): *Rabbit erythrocyte haemolysis assay for α -toxin*. The rabbit red blood cell (RBC) haemolysis assay was used to determine biologically-relevant levels of α -toxin released during *S. aureus* culture^{19,20}. Fresh, heparinised RBCs (Orygen, Penicuik, UK), were prepared at $\sim 5\%$ haematocrit in Hank's balanced salt solution (HBSS; pH 7.4; Invitrogen, Paisley, UK). At time zero and at specific points throughout the culture of *S. aureus* (see Figures), aliquots were taken and centrifuged (8,000xg; 30secs) and the supernatant aliquotted into microcentrifuge tubes, taking care not to disturb the pellet. The supernatants which contained α -toxin, were

154 then added to microcentrifuge tubes containing 5% RBC suspensions, mixed gently and incubated
 155 (37°C;60mins). The samples were then centrifuged (8,000xg;10secs) and the supernatant assessed for
 156 haemoglobin at 540nm (Abs₅₄₀) on a Nanodrop spectrophotometer (ThermoFisher Scientific, Waltham, USA).
 157 Percent haemolysis (%H) was then calculated ((Abs₅₄₀ of sample – Abs₅₄₀ of negative control)/(Abs₅₄₀ of 100%
 158 haemolysis – Abs₅₄₀ of negative control) x100%). The sensitivity of RBC samples to α -toxin and the amount
 159 of α -toxin produced from each culture were variable. It was important to minimise this at the beginning of an
 160 experimental week, by initially testing erythrocytes from several rabbits on a freshly-prepared batch of α -toxin.
 161 Erythrocytes which were relatively insensitive to α -toxin requiring >1hr of incubation before haemolysis was
 162 detectable, were not studied further and this reduced the variation in results obtained.

163 *(d): Assessment of in situ chondrocyte viability by confocal scanning laser microscopy (CLSM).* Fresh, healthy
 164 (non-degenerate) metacarpophalangeal joints of 3-yr old cows from a local abattoir, were washed, skinned,
 165 de-hoofed and opened under aseptic conditions. Cartilage explants were cultured in Dulbecco's Modified
 166 Eagle's medium (DMEM;pH 7.4;37°C; Invitrogen) in the presence or absence of *S. aureus* and after 15hrs the
 167 bacteria had caused measurable (~20%) chondrocyte death. Explants were then exposed to antibiotics (or
 168 control medium) and the incubation continued for up to 23hrs. At the time points indicated, explants were
 169 removed, rinsed in DMEM and chondrocyte viability assessed as described³⁰. Explants were incubated with 5-
 170 chloromethylfluorescein diacetate (CMFDA) and propidium iodide (PI; both at 10 μ M from Invitrogen, Paisley,
 171 UK) in DMEM (1hr;21°C). These probes labelled living or dead cells green or red, respectively. Explants were
 172 then fixed (4% formaldehyde;4hrs;Fisher Scientific, Loughborough, UK), rinsed and stored in PBS, until
 173 imaged within 24hrs. Imaging of fluorescently-labelled *in situ* chondrocytes was performed as described^{14,30}.
 174 An upright Zeiss LSM510 Axioskop (Carl Zeiss, Welwyn Garden City, UK) CLSM with a x10 (0.3NA) dry
 175 objective was used to acquire axial images. The percentage cell death ((number of dead cells/total number of
 176 living and dead cells) x100%) was calculated within a standard region of interest (ROI) using VolocityTM 4
 177 software (Improvision, Coventry, UK). Within each ROI, individual cells, in both green (living) and red (dead)
 178 channels, were quantified based on percentage voxel (volumetric pixel) intensity¹⁴.

179 *(e): Data analysis and statistics.* Data are shown as means \pm standard error of the mean (S.E.M) from at least
 180 3 independent experiments, and were analysed using GraphPad Prism Ver.6 (GraphPad, San Diego, CA, USA).
 181 Non-parametric t-tests (Kruskal-Wallis) and ANOVAs were performed and significant differences accepted
 182 when $P<0.05$.

183

Results

(a) *S. aureus* α -toxin mediated rabbit RBC haemolysis. The strain of *S. aureus* (DU5946) that harboured the α -toxin (*hla*) gene was haemolytic on rabbit blood agar whereas the strain deficient in this gene (DU5938) was non-haemolytic (Fig. 1(a)). Supernatants were prepared from cultures of these strains and added to suspensions of rabbit erythrocytes in HBSS. There was a rapid increase in %H for the α -toxin-positive strain, however the α -toxin-deficient strain had no detectable effect (Fig. 1(b)). Thus, by using rabbit erythrocytes in suspension as a model and determining haemoglobin release spectrophotometrically, biologically active α -toxin could be quickly assessed semi-quantitatively.

(b) *Time course of linezolid action.* Single doses of linezolid (to concentrations of 5-50 μ g/ml) were added to the α -positive *S. aureus* cultures (DU5946) at a point corresponding to ~50% haemolysis (after 80mins) and samples analysed for α -toxin-induced haemolysis (Fig. 2(a)). For all concentrations, there was a rapid reduction in the rate of rise of α -toxin-induced haemolysis, and at 25 and 50 μ g/ml, there was no significant change in %H (Fig. 2(b)). Addition of linezolid to 50 μ g/ml at 80, 100, 130mins time points, rapidly suppressed any further increase in α -toxin-induced haemolysis (Fig. 2(c)). Linezolid (50 μ g/ml) addition 60mins after the commencement of the *S. aureus* culture, abolished subsequent α -toxin-induced haemolysis compared to the untreated (infected) control (Fig. 3(a)). The same concentration of linezolid added after 100mins when there was ~40% haemolysis, also completely inhibited subsequent α -toxin-induced haemolysis to the same level after 40 and 60mins as that present at the start of the experiment ($P<0.001$; Fig. 3(b)). These results indicated that the linezolid addition to *S. aureus* cultures rapidly (within ~20 mins) inhibited α -toxin-induced haemolysis and that 25 and 50 μ g/ml prevented any further rise.

(c) *Time course of penicillin action.* Single doses of penicillin G (to 5000, 10,000, 25,000 or 50,000U/ml) were added to the α -toxin positive *S. aureus* cultures at a time point corresponding to ~50% haemolysis (after 60mins) and samples analysed for α -toxin-induced haemolysis (Fig. 4(a)). While at penicillin concentrations of 50,000U/ml and 250,000U/ml at 140 mins there was a significant reduction in the rate of rise of α -toxin-induced haemolysis, the effect of lower concentrations was not significant (Fig. 4(b)). Addition of penicillin to 50,000U/ml at different time points, suppressed the increase in α -toxin-induced haemolysis (Fig. 4(c)), however the time course of action was slower compared to linezolid (Fig. 2(c)). Penicillin addition 20mins after the start of the culture, significantly inhibited haemolysis compared to the untreated (infected) control after 40 and 60mins (Fig. 5(a)). While in the presence of penicillin, there were significant increases in haemolysis at 40 and 60mins compared to the values at the time of drug addition ($P<0.05$), these were not different ($P=0.12$), indicating a plateau in α -toxin levels. When the same concentration of penicillin was added after 60mins, there was ~60% haemolysis at time zero and this significantly ($P<0.001$) inhibited subsequent α -toxin-induced haemolysis after 40 and 60mins (Fig. 5(b)). The extent of α -toxin-induced haemolysis at 60mins was significantly ($P=0.05$) higher than the time zero point but not significantly different compared to the 40min time point (Fig. 5(b)). These results suggested that penicillin addition to *S. aureus* cultures progressively reduced the rate of increase of α -toxin-induced haemolysis. However, the time course appeared

222 relatively slow compared to linezolid, and at the highest concentrations studied (50,000U/ml) there was still a
223 slight rise in α -toxin-induced haemolysis until the inhibition appeared complete.

224 To directly compare the inhibitory effects of linezolid and penicillin using the same blood samples and
225 *S. aureus* cultures, DU5946 cultures were incubated for 60mins, and then untreated (control) or dosed with
226 penicillin (250,000U/ml) or linezolid (25 μ g/ml), and the medium analysed for α -toxin for up to 80mins (Fig.
227 6). After 20mins, both antibiotics significantly ($P<0.05$) inhibited haemolysis, however at 40, 60 and 80mins
228 after antibiotic addition, the inhibition was significantly (by $\sim 8\%$) greater ($P<0.05$) in the presence of linezolid
229 (Fig. 6). The inhibition by penicillin did not change significantly compared to linezolid over the 40-80mins
230 period after antibiotic addition, suggesting that the inhibitory effect of penicillin was maximal.

231 *(d) Chondrocyte protection in a bovine cartilage septic arthritis model by antibiotics.* *S. aureus* DU5946 was
232 cultured with cartilage explants and in control (no antibiotics) samples, there was almost complete ($>90\%$)
233 chondrocyte death after ~ 21 hrs (Fig. 7(a,b)). In contrast, when infected cartilage was treated with penicillin or
234 linezolid after 15hrs, there was no subsequent chondrocyte death and no difference between the potency of the
235 two drugs (Fig. 7(a,b)). Similar experiments were performed with flucloxacillin and erythromycin (data not
236 shown). In the untreated infected samples, there was complete ($\sim 100\%$) chondrocyte death after ~ 24 hrs.
237 Treatment of infected cartilage samples with flucloxacillin (2mg/ml) or erythromycin (10 μ g/ml) after 18hrs,
238 prevented any subsequent chondrocyte death and there was no significant difference ($P>0.05$) between the two
239 drugs at any time point studied up to 26hrs after antibiotic addition. Levels of chondrocyte death in the control
240 (i.e. uninfected and untreated) cartilage samples were $<2\%$. These results suggested that both bacteriolytic
241 (penicillin, flucloxacillin) and bacteriostatic/bactericidal (linezolid, erythromycin) antibiotics were equally
242 effective at protecting chondrocytes in the bovine cartilage model of *S. aureus* septic arthritis.

243

Discussion

The rabbit erythrocyte haemolysis assay determined the time course of biologically-relevant α -toxin produced during culture of *S. aureus* in the presence or absence of bacteriostatic/bactericidal and bacteriolytic antibiotics. The antibiotics tested could limit or prevent a rise in α -toxin levels, however there were differences in their time courses of action. The bacteriostatic/bactericidal agent, linezolid, appeared more rapid and potent than the bacteriolytic antibiotic, penicillin. There was, however, no evidence of a detectable 'pulse' of α -toxin-induced haemolysis following addition of any of the antibiotics studied, suggesting there was no substantial intracellular 'pool' of stored α -toxin in the *S. aureus* strain DU5946. In the bovine cartilage model of septic arthritis, *S. aureus* cultures rapidly caused chondrocyte death. However chondrocytes were completely protected against *S. aureus* by the antibiotics tested (linezolid, flucloxacillin, penicillin, erythromycin). These results were important for elucidating whether or not there was a significant intracellular store of α -toxin as this may be relevant for antibiotic selection during *S. aureus*-induced septic arthritis.

The rabbit RBC haemolysis bio-assay was used to determine the time course of α -toxin release during *S. aureus* culture. The measurement of haemoglobin released following the interaction between α -toxin and erythrocytes provided a dynamic and sensitive assay for the lethal effects of α -toxin, and was highly flexible experimentally and reproducible, allowing several conditions (e.g. drug concentrations) to be tested at the same time points. The other toxins/damaging agents produced by *S. aureus* had a negligible effect on rabbit erythrocytes emphasising the specific nature of this assay for α -toxin (Fig. 1). It was noted that some blood samples were very sensitive to α -toxin whereas others were relatively unresponsive and so it was necessary to initially screen each blood sample for sensitivity against a *S. aureus* culture. Once a blood sample was deemed acceptable, the time course of haemolysis to α -toxin was highly reproducible. In addition, the ability of *S. aureus* cultures to produce sufficient quantities of potent α -toxin was variable and probably depended on prevailing culture conditions. The difference in potency of batches of α -toxin was observed, as for example, in one series of experiments, 60mins elapsed before the *S. aureus* culture started to produce measurable α -toxin levels (Fig. 2(a)). In another set of experiments performed with a different α -toxin preparation, after only 40mins haemolysis had already increased to 20% (Fig. 4(a)). The rabbit erythrocyte model for determining the effects of antibacterials on α -toxin levels might be considered a limitation. However, rabbit erythrocytes contain the *S. aureus* α -toxin receptor ADAM-10 also present on chondrocytes of animals and normal and degenerate human cartilage¹⁶⁻¹⁸. Our previous work has demonstrated that in a cartilage model of septic arthritis, α -toxin is the key damaging agent to chondrocytes and thus the rabbit erythrocyte hemolysis assay is appropriate to determine the effects of antibacterials on α -toxin release from *S. aureus* cultures^{14,15}.

Linezolid at 25 or 50 μ g/ml rapidly abolished the rise in α -toxin-induced haemolysis (Fig. 2(c)) probably because of its potent action on bacterial protein synthesis and replication³¹. However, linezolid may have both bacteriostatic and/or bactericidal effects on *S. aureus* depending on concentration and experimental conditions²². Therefore the reduced rate of haemolysis in linezolid-treated cultures may be the result of bacteriostatic, bactericidal or combined bacteriostatic/bactericidal action. Whatever the mechanism, at 50 μ g/ml linezolid (Fig. 3(a,b)), there was no progressive change in haemolysis ((Fig. 2(c); Fig. 3(b)) indicating no additional release of α -toxin from dying/dead bacteria. However, future work would be required to

investigate the effect of varying concentrations of linezolid on the bacteriostatic and/or bactericidal modes of actions on *S. aureus* α -toxin. It should be noted that to test our hypothesis of whether there was a detectable intracellular store of α -toxin, we used concentrations considerably above the MIC (minimum inhibitory concentrations; Table 2) and have not attempted to study the effects of clinically-relevant doses.

The action of penicillin was slower compared to linezolid as evidenced by the relatively tardy inhibition of α -toxin-induced haemolysis even at a high concentration (50,000U/ml; Fig. 4(c); Fig. 5(a,b)). There are two possible explanation for this finding. The most likely interpretation of these results is because of the slower time course of the antibiotic inhibiting bacterial cell wall synthesis^{21,22}. The level of haemolysis after penicillin addition was higher than that with linezolid, and did not change with time (Fig. 6). This suggested that the inhibition by both drugs was maximal with no further bacterial death or inhibition of division. However it is also possible that the slightly elevated level of haemolysis (~8% of the total) with a high dose of penicillin (Fig. 6) could have been because of the release of the small additional amount of intracellular α -toxin from the dying/dead bacteria.

The presence of a small amount of intracellular *S. aureus* α -toxin is implied by others using different methods to cause bacterial damage. For example Duncan and Cho³² suggested that mechanical disruption of *S. aureus* released 1-2% of total toxin present in cultures. However this might have been influenced by the simultaneous release of degradative enzymes leading to an underestimation in the amount of active toxin. Monecke et al³³ using alkali treatment, noted that there was only a 'small' (but unquantified) level of stored α -toxin following bacterial lysis. For *S. aureus* therefore, there might only be a small intracellular store with the α -toxin released rapidly after synthesis. On the other hand, pneumolysin, the toxin produced by *Streptococcus pneumoniae*, is almost exclusively stored intra-cellularly and only released through bacterial autolysis^{34,35}. In a *S. pneumoniae* rabbit meningitis model, addition of the bacteriolytic agent ceftriaxone, released substantial additional quantities of pneumolysin compared to the non-bacteriolytic agent rifampin, with the additional toxin causing further cellular damage and inflammation³⁶.

In the *S. aureus* septic arthritis cartilage model, linezolid and penicillin (Fig. 7a,b) and flucloxacillin and erythromycin (see Results) were equally effective. This might appear to conflict with the suggestion above that there was a small intracellular 'pool' of α -toxin. It is possible that the methods for determining chondrocyte death were not sufficiently sensitive to detect the small increase from the release of stored α -toxin by penicillin. However there was no significant difference in % chondrocyte viability between linezolid and penicillin-treated explants so this does not seem likely. Alternatively, α -toxin levels in the medium and thus the cartilage matrix may already be high and any small increase following penicillin treatment could be without further detectable effect. The α -toxin released from *S. aureus* will have a rapid effect on the rabbit erythrocytes as pore formation and haemolysis proceeds promptly¹⁹. However in cartilage, toxin permeability may be hindered by the cartilage extracellular matrix³⁷. It is also possible that the dynamics of toxin binding and pore formation to bovine chondrocytes proceeds with a different time course compared to rabbit erythrocytes. ADAM-10 (A Disintegrin And Metalloproteinase domain-containing protein-10) has been identified as an important receptor for α -toxin binding³⁸ mediating changes to intracellular Ca^{2+} signalling³⁹ and is present on human chondrocytes

320 and up-regulated in osteoarthritis¹⁶⁻¹⁸. The binding site density on chondrocytes may also be different compared
321 to rabbit erythrocytes which could further influence the time course of the α -toxin effect.

322 The rapid inhibition by linezolid on the rise in *S. aureus* α -toxin levels compared to penicillin is likely
323 due to a bacteriostatic action on *S. aureus* replication, and *in vivo* this would be followed by bacteria removal
324 by the immune system. Septic arthritis may be regarded as a medical emergency³ and thus clinically there
325 would be benefit in administering this antibacterial agent or others in this class for the most rapid action
326 possible to protect joint tissues from further damage. It is possible that bacteriolytic antibiotics (e.g. penicillin,
327 flucloxacillin) may lead to the release of additional inflammatory components (e.g. teichoic acids,
328 peptidoglycans) from *S. aureus* potentially causing further chondrocyte damage. However all the drugs tested
329 (penicillin, linezolid (Fig. 7); flucloxacillin, erythromycin (see Results)), were equally effective at protecting
330 chondrocytes against *S. aureus*. This suggests that these agents, if released from dead/dying bacteria, do not
331 have a significant effect on chondrocyte viability and support the view that the α -toxin alone was the primary
332 cause of chondrocyte death¹⁵. It is also worth noting that bacteriostatic/bactericidal agents e.g. linezolid, reduce
333 the expression of *S. aureus* α -toxin^{29,31}. This is in marked contrast to sub-inhibitory doses of β -lactams which
334 increase *S. aureus* hla (α -toxin) mRNA expression^{40,41} and therefore might cause concern in the treatment of
335 osteoarticular infection by these bacteria. Furthermore, linezolid reduces expression of other staphylococcal
336 toxins such as PVL in PVL-associated staphylococcal pneumonia⁴². Since PVL-producing strains of *S. aureus*
337 may also be present and can cause complications in septic arthritis⁴³, linezolid may offer a further advantage
338 over bacteriolytic antibiotics. Although our results demonstrated that these antibacterials are protective against
339 chondrocytes *in vitro*, the response *in vivo* is likely to be far more complex as our model of *S. aureus*-induced
340 septic arthritis does not include the host's immune response.

341 While it can be particularly challenging to extrapolate the results from the *in vitro* situation to the
342 clinical setting, there are three main findings from this study which are relevant. (a) The time course of the
343 bacteriostatic/bactericidal antibiotic linezolid on limiting the rise α -toxin was clearly more rapid than for the
344 bacteriolytic antibiotic penicillin (Fig. 2 vs Fig. 4). (b) There was a negligible amount of stored α -toxin released
345 following antibacterial treatment (Fig. 6). (c) There was no significant difference in the chondroprotective
346 effect between exemplars of the two classes of antimicrobials in the septic arthritis model (Fig. 7). It is also
347 relevant to note that Monecke et al.³³ have presented evidence to indicate that secretion of *S. aureus* α -toxin
348 does not appear to correlate with the progression of septic arthritis. There is nevertheless a strong link between
349 levels of α -toxin and chondrocyte death^{14,15} and it is known that cartilage which does not possess viable
350 chondrocytes will degenerate⁴⁴. Thus, while there is probably a relationship between levels of α -toxin in *S.*
351 *aureus* – induced septic arthritis and *in vivo* cartilage viability, a direct correlation is yet to be established.
352 Despite this, the dominant importance of α -toxin produced by *S. aureus* in septic arthritis as opposed to other
353 damaging agents released by this bacterium, has been underscored by recent preliminary *in vivo* studies⁴⁵. In
354 these experiments, the intra-articular injection of *S. aureus* 8325-4 which produces all toxins including α -toxin
355 and other damaging agents, was compared to the injection of an *S. aureus* mutant (DU1090) which produces
356 the full range of toxins/enzymes, *except* α -toxin¹⁵. Mice injected with either *S. aureus* strains developed septic
357 arthritis with evidence of weight loss, limb swelling and gait changes whereas these were absent in the control

358 (PBS – injected) group. Notably, there was significantly ($P<0.05$) more chondrocyte death in the group infected
359 with *S. aureus* 8325-4 (approx. 93% chondrocyte death) when compared to DU1090 (approx. 26% chondrocyte
360 death) and PBS-injected (i.e. control) mice (5% chondrocyte death). The results suggested that α -toxin was the
361 major chondrocyte damaging agent, but also any adverse effect of the immune system during this time course
362 with this animal model of septic arthritis was negligible in comparison. While antimicrobial treatment of *S.*
363 *aureus* to limit further production of α -toxin is clearly an essential clinical approach, strategies aimed at quickly
364 suppressing the action of α -toxin already present in the infected tissue could be of additional benefit. For
365 example, the damaging effect of α -toxin is markedly suppressed by raising the osmolarity of the culture
366 medium²⁰.

367 This present study utilised a rabbit RBC assay to determine α -toxin levels produced by *S. aureus* in
368 culture following treatment with antibiotics. Linezolid rapidly limited any further increase in α -toxin levels by
369 *S. aureus* whilst penicillin had a slower time course of action. There was a small additional release of α -toxin
370 following penicillin addition suggesting a delayed action and/or a small intracellular store of α -toxin. However,
371 both drugs were equally effective at protecting chondrocytes in our *S. aureus* septic arthritis model.

372

373 **Acknowledgements.** We thank Prof. T.J. Foster, Department of Microbiology, Trinity College, Dublin, for
374 supplying *S. aureus* strains, Dr A. Kubasik-Thayil IMPACT facility, University of Edinburgh, for CLSM
375 guidance and K. Wójcik of Scotbeef Ltd, Bridge of Allan, Stirlingshire, for providing bovine feet. We also
376 thank A. Boyle and Dr C.J. Doherty for helpful comments. This work was supported by Pfizer Grant No:
377 WI177684.

378

379 **Conflict of Interest Statement:** The authors do not have any conflicts of interest to declare in relation to the
380 work presented in this study.

381

References

1. Mathews CJ, Kingsley G, Field M, Jones A, Weston V *et al.* Management of septic arthritis: a systematic review. *Postgrad Med J* 2008;84(991):265-270.
2. Kaandorp CJ, Krijnen P, Moens HJ, Habbema JD, van Schaardenburg D. The outcome of bacterial arthritis: a prospective community-based study. *Arthritis Rheum* 1997;40:884-892.
3. Goldenberg DL. Septic arthritis. *Lancet* 1998;351:197-202.
4. Krieg AM. A possible cause of joint destruction in septic arthritis. *Arthritis Res Ther* 1999;1:3-4.
5. Mathews CJ, Weston VC, Jones A, Field M, Coakley G. Bacterial septic arthritis in adults. *Lancet* 2010;375:846-855.
6. Wang DA, Tambyah PA. Septic arthritis in immunocompetent and immunosuppressed hosts. *Best Pract Res Clin Rheumatol* 2015;29:275-289.
7. Balato G, Di Donato SL, Ascione T, D'Addona A, Smeraglia F *et al.* Knee septic arthritis after arthroscopy: incidence, risk factors, functional outcome and infection eradication rate. *Joints* 2017;7(2):107-113.
8. Gupta MN, Sturrock RD, Field M. A prospective 2-year study of 75 patients with adult-onset septic arthritis. *Rheumatology (Oxford)* 2001;40:24-30.
9. Sampedro GR, Bubeck Wardenburg J. *Staphylococcus aureus* in the intensive care unit: are these golden grapes ripe for a new approach? *J Infect Dis* 2017;215(S1):S64-S70.
10. Muir H. The chondrocyte, architect of cartilage. Biomechanics, structure, function and molecular biology of cartilage matrix macromolecules. *Bioessays* 1995;17:1039-1048.
11. Shirtliff ME, Mader JT. Acute septic arthritis. *Clin Microbiol Rev* 2002;15:527-544.
12. Powers ME, Bubeck Wardenburg J. Igniting the fire: *Staphylococcus aureus* virulence factors in the pathogenesis of sepsis. *PLoS Pathog* 2014;10(2):e1003871. doi: 10.1371/journal.ppat.1003871.
13. Dinges M, Orwin P, Schlievert P. Exotoxins of *Staphylococcus aureus*. *Clinical Microbiol Rev* 2000;13(1):16-34.
14. Smith IDM, Winstanley JP, Milto KM, Doherty CJ, Czarniak E *et al.* Rapid chondrocyte death induced by *Staphylococcus aureus* toxins in a bovine cartilage explant model of septic arthritis. *Osteoarth Cart* 2013;21(11):1755-1765.
15. Smith IDM, Milto KM, Doherty CJ, Amyes SGB, Simpson AHRW *et al.* A potential key role for alpha-haemolysin of *Staphylococcus aureus* in mediating chondrocyte death in septic arthritis. *Bone Jt Res* 2018;7(7):457-467. doi: 10.1302/2046-3758.77.BJR-2017-0165.R1
16. Chubinskaya S, Cs-Szabo G, Kuettner KE. ADAM-10 message is expressed in human articular cartilage. *J. Histochem. Cytochem.* 1998;46(6):723-729.
17. Chubinskaya S, Mikhail R, Deutsch A, Tindal MH. ADAM-10 protein is present in human articular cartilage primarily in the membrane-bound form and is upregulated in osteoarthritis and in response to IL-1 α in bovine nasal cartilage. *J. Histochem. Cytochem.* 2001;49:1165-1176.
18. Wilke GA, Bubeck Wardenburg J. Role of a disintegrin and metalloprotease 10 in *Staphylococcus*

aureus alpha-hemolysin-mediated cellular injury. *Proc Natl Acad Sci USA* 2010;107(30):13473-13478.

19. Cooper LZ, Madoff MA, Weinstein L. Hemolysis of rabbit erythrocytes by purified Staphylococcal alpha-toxin. I. Kinetics of the lytic reaction. *J Bacteriol* 1964;87:127-135.
20. Liu, C.L.S. & Hall, A.C. Optimising the composition of irrigation fluid to reduce the potency of *S. aureus* α -toxin: a possible role in the treatment of septic arthritis. *Cartilage* 2018; IN PRESS <https://doi.org/10.1177/1947603518798888>
21. Bergeron MG, Bruschi JL, Barza M, Weinstein L. Bactericidal activity and pharmacology of flucloxacillin. *Amer J Med Sci* 1976;271(1):13-20.
22. Elliott TS, Greenwood D, Rodgers FG, O'Grady F. The response of *Staphylococcus aureus* to benzylpenicillin. *Br J Exp Pathol* 1979;60(1):14-23.
23. Hobson D. Activity of erythromycin against *Staphylococcus aureus*. *Br Med J* 1954;30.1(4856):236-239.
24. Le Noc P, Croize J, Bryskier A, Le Noc D, Robert J. Comparative in vitro bacteriostatic and bactericidal effect of 5 macrolides: roxithromycin, erythromycin, oleandomycin, josamycin and spiramycin against 284 hospital bacterial strains. *Pathol Biol (Paris)* 1989;37(5 Pt 2):553-559
25. Brauers J, Kresken M, Menke A, Orland A, Weiher H *et al.* Bactericidal activity of daptomycin, vancomycin, teicoplanin and linezolid against *Staphylococcus aureus*, *Enterococcus faecalis* and *Enterococcus faecium* using human peak free serum drug concentrations. *Int J Antimicrob Agents* 2007;29:322-325.
26. Nilsson I, Hartford O, Foster T, Tarkowski N. Alpha-toxin and gamma-toxin jointly promote *Staphylococcus aureus* virulence in murine septic arthritis. *Infect Immun* 1999;67:1045-1049.
27. Herbert S, Ziebandt AK, Ohlsen K, Schafer T, Hecker M *et al.* Repair of global regulators in *Staphylococcus aureus* 8325 and comparative analysis with other clinical isolates. *Infect Immun* 2010;78:2877-2889.
28. Novick R. Properties of a cryptic high-frequency transducing phage in *Staphylococcus aureus*. *Virology* 1967;33:155-166.
29. Gemmell C, Goutcher S, Reid R, Sturrock R. Role of certain virulence factors in a murine model of *Staphylococcus aureus* arthritis. *J Med Microbiol* 1997;46:208-213.
30. Amin AK, Bush PG, Huntley JS, Simpson AHW, Hall AC. Osmolarity influences chondrocyte death in wounded articular cartilage. *J Bone Jt Surg (Amer)* 2008;90:1531-1542.
31. Bernardo K, Pakulat, N, Fleer S, Schnaith A, Utermöhlen O *et al.* Subinhibitory concentrations of linezolid reduce *Staphylococcus aureus* virulence factor expression. *Antimicrob Agents Chemother* 2004;48(2):546-555.
32. Duncan JL, Cho GJ. Production of Staphylococcal alpha toxin. I Relationship between cell growth and toxin formation. *Infect. Immun.* 1971;4(4):456-461.
33. Monecke S, Muller E, Buchler J, Stieber B, Ehrlich R. *Staphylococcus aureus* in vitro secretion of alpha toxin (hla) correlates with the affiliation to clonal complexes. *PLoS One* 2014;9(6):e100427.
34. Berry AM, Lock RA, Hansman D, Paton JC. Contribution of autolysin to virulence of *Streptococcus pneumoniae*. *Infect Immun* 1989;57:2324-2330.

35. Mitchell AM, Mitchell TJ. *Streptococcus pneumoniae*: virulence factors and variation. *Europ Soc Clin Micro Infect Dis* 2010;16:411-418.
36. Spreer A, Kerstan H, Bottcher T, Gerber J, Siemer A *et al*. Reduced release of pneumolysin by *Streptococcus pneumoniae* *in vitro* and *in vivo* after treatment with nonbacteriolytic antibiotics in comparison to ceftriaxone. *Antimicrob Agents Chemother* 2003;47(8), 2649-2654.
37. Maroudas A. Physico-chemical properties of articular cartilage. In: Freeman, MAR, ed. *Adult Articular Cartilage*. London: Pitman Medical. 1979. pp. 215-290.
38. Von Hoven G, Rivas AJ, Neukirch C, Klein S, Hamm C *et al*. Dissecting the role of ADA-10 as a mediator of *Staphylococcus aureus* α -toxin action. *Biochem J* 2016;473(13):1929-40. doi: 10.1042/BCJ20160062
39. Seilie ES, Bubeck Wardenburg J. *Staphylococcus aureus* pore-forming toxins: The interface of pathogen and host complexity. *Sem Cell Dev Biol* 2017;72:101–116.
40. Stevens DL, Ma Y, Salmi DB, McIndoo E, Wallace RJ *et al*. Impact of antibiotics on expression of virulence-associated exotoxin genes in methicillin-sensitive and methicillin-resistant *Staphylococcus aureus*. *J Infect Dis* 2007;195:202-211.
41. Kernodle DS, McGraw PA, Barg NL, Menzies BE, Voladri RK, Harshman S. Growth of *Staphylococcus aureus* with nafcillin *in vitro* induces alpha-toxin production and increases the lethal activity of sterile broth filtrates in a murine model. *J. Infect. Diseases*. 1995;172(2):410-419.
42. Morgan MS. Diagnosis and treatment of Panton-Valentine leukocidin (PVL)-associated staphylococcal pneumonia. *Int J Antimicrob Agents* 2007;30:2289-2296.
43. Wright JA, Nair SP. Interaction of staphylococci with bone. *Int J Med Microbiol* 2010;300:193-204.
44. Paterson, S.I., Eltawil, N.M., Simpson, A.H.R.W., Amin, A.K. & Hall, A.C. (2016). Drying of open animal joints *in vivo* subsequently causes cartilage degeneration. *Bone Jt. Res.* 5:137-144.
45. Clement, R., Hall, A.C., Howie, S.E.M. Simpson, A.H.R.W. (2016). A murine model of septic arthritis demonstrates that infection with an alpha toxin-producing strain of *S. aureus* leads to significantly elevated levels of chondrocyte death within 48 hours of infection when compared with infection with an alpha toxin-deficient mutant strain. *Bone Joint J.* 98-B (SUPP 16): 20.
46. Andrews JM. Determination of minimum inhibitory concentrations. *J Antimicrob Chemother* 2001;48:Suppl. 1: 5-16.
47. Gemmell CG. Virulence factor expression by Gram-positive cocci exposed to subinhibitory concentrations of linezolid. *J Antimicrob Chemother* 2002;50(5):665-672.
48. Gemmell CG. Susceptibility of a variety of clinical isolates to linezolid: a European inter-country comparison. *J Antimicrobial Chemother* 2001;48(1):47-52.

Tables

Table 1. Miller et al., (2018)

<i>S. aureus</i> strain	Genotype	Phenotype	Haemolysins produced	Given name
DU5946	<i>Hlb::φ42E Δhlg::Tc^r</i>	Hla ⁺ Hlb ⁻ Hlg ⁻	Hla	α-toxin positive α ⁺ β ⁻ γ ⁻
DU5938	<i>Hlb::φ42E hla::Em^r Δhlg::Tc^r</i>	Hla ⁻ Hlb ⁻ Hlg ⁻	None	α-toxin deficient α ⁻ β ⁻ γ ⁻

Table 1. The bacterial strains used in study. The strain of *S. aureus* is shown with its genotype, phenotype and the toxins produced. Tc^r = Tetracycline resistance. Em^r = Erythromycin resistance.

Antibiotic	Class (action)	Mode of action (on bacteria)	Antibiotic Dose		
			Dose used	MIC	Reference
Linezolid	Oxazolidinone (bacteriostatic/ bactericidal)	Ribosomal 50S inhibitor – interrupts protein synthesis	50µg/ml for haemolysis experiments; 25µg/ml for cartilage experiments.	0.12-8µg/ml	46
Erythromycin	Macrolide (bacteriostatic/ bactericidal)	Ribosomal 50S inhibitor – interrupts protein synthesis	10µg/ml for cartilage experiments.	0.06-128µg/ml	46
Penicillin	β-lactam (bacteriolytic)	Blocks cross- linking of peptidoglycan - inhibits cell wall formation	50,000U/ml (34mg/ml) for haemolysis experiments; 1000U/ml (0.70µg/ml) for cartilage experiments.	0.015-128µg/ml	46
Flucloxacillin	β-lactam (bacteriolytic)	Blocks cross- linking of peptidoglycan - inhibits cell wall formation	2mg/ml for cartilage experiments.	0.4µg/ml (fully effective)	21

552 **Table 2. Antibiotics and concentrations used in study.** The class, mode of action and concentration of
553 antibiotics used are shown. The MIC (minimum inhibitory concentration) for linezolid against *S. aureus* ranges
554 over 0.12-8µg/ml^{46,47}; and the MIC90 (minimum inhibitory concentration required to block 90% bacterial
555 growth) ranges over 2-4µg/ml⁴⁸. Doses used in the present work ranged from 5-50µg/ml and were effective
556 (Fig. 2(c), Fig. 7a)). The MIC for erythromycin on Staphylococci ranges over 0.06-128µg/ml⁴⁶ and the dose
557 used for cartilage experiments was 10µg/ml. The MIC for penicillin action on Staphylococci ranges over 0.015-
558 128µg/ml⁴⁶. The present study used 50,000U/ml (34mg/ml) for haemolysis experiments and 1000U/ml
559 (0.70µg/ml) for cartilage experiments which were effective (Figs. 4(c) and 7); see text). Flucloxacillin has
560 been shown to be completely effective against *S. aureus* when present at 0.4µg/ml²¹. In the present study a
561 concentration of 2mg/ml was used (see text).

Figure Legends

Figure 1. The haemolysis of rabbit erythrocytes following incubation with α -toxin positive or α -toxin negative strains of *S. aureus*. (a) Cultures of *S. aureus* (i) α -toxin positive, β and γ -toxin negative strain or (ii) α -toxin negative, β and γ -toxin negative strain (DU5938; Table 1) were incubated on tryptone soya agar (TSA) containing rabbit erythrocytes. Clear areas surrounding colonies of α -toxin secreting *S. aureus* represent the haemolysis of rabbit erythrocytes. No haemolysis was observed around colonies of the α -toxin negative strain. (b) The time course of haemolysis in suspensions of rabbit erythrocytes during culture with α -toxin positive or α -toxin negative strain of *S. aureus*. For this and subsequent Figures, when error bars were not shown they were smaller than the symbols. Data shown are means \pm s.e.m. for 3 independent experiments for each condition.

Figure 2. Effects of linezolid on α -toxin-induced haemolysis of rabbit erythrocytes. Panel (a) shows the effect of linezolid (0 (HBSS control), 5, 10, 25 or 50 μ g/ml) addition at 80mins to *S. aureus* α -toxin positive strain (DU5946) cultures on α -toxin-induced haemolysis (see Materials and Methods for details). Panel (b) presents the rate of change of haemolysis over 100-140mins ((expressed as the % change/min) \times 10) extent following addition of linezolid with increasing concentration. Asterisks (* or **) represent significant differences at the $P < 0.05$ or $P < 0.001$ levels respectively. Panel (c) demonstrates the effects of linezolid (50 μ g/ml) addition to *S. aureus* cultures at 60, 80, 100 and 130mins corresponding to approximately 0, 10, 40 or 80% α -toxin-induced haemolysis. Data shown are means \pm s.e.m. for at least 3 independent experiments.

Figure 3. Inhibition of α -toxin-induced haemolysis of rabbit erythrocytes at different time points by prior treatment with linezolid. Linezolid (50 μ g/ml) was added (a) 60mins or (b) 100mins after the initiation of *S. aureus* α -toxin positive strain (DU5946) cultures which corresponded to approximately 0 or 40% α -toxin-induced haemolysis respectively at the start of the experiment. Measurements of %H were then taken at 0, 40 and 60mins as shown. At 40 and 60mins, the prior treatment with linezolid significantly ($P < 0.001$) reduced α -toxin-induced haemolysis to levels that were not significantly different from the control (i.e. levels of %H at the start of the time course). Data shown are means \pm s.e.m. for at least 3 independent experiments.

Figure 4. Effects of penicillin on α -toxin-induced haemolysis of rabbit erythrocytes. Panel (a) shows the effect of penicillin (0, 5,000, 10,000, 25,000 or 50,000U/ml) addition at 60mins to α -toxin positive strain (DU5946) *S. aureus* cultures on α -toxin-induced haemolysis (see Materials and Methods for details). Panel (b) presents the extent of α -toxin-induced haemolysis at 140mins following addition of penicillin. The inhibition by penicillin was only significantly different from the HBSS control at 50,000 and 250,000U/ml ($P < 0.001$) whereas there was no significant difference between these two antibiotic concentrations. Panel (c) demonstrates the effects of penicillin (50,000 U/ml) addition to *S. aureus* cultures at 20, 40, 60 and 100 mins

599 corresponding to approximately 0, 10, 40 or 80% α -toxin-induced haemolysis. Data shown are means \pm s.e.m.
600 for at least 3 independent experiments.

601

602 **Figure 5. Inhibition of α -toxin-induced haemolysis of rabbit erythrocytes at different time points by**
603 **prior treatment with penicillin.** Penicillin (50,000U/ml) was added (a) 20mins or (b) 60mins after the
604 initiation of α -toxin positive strain (DU5946) *S. aureus* cultures which corresponded to approximately 0 or
605 60% α -toxin-induced haemolysis at the start of the experiment. Measurements of %H were then taken at 0, 40
606 and 60mins as shown. At 40 and 60mins, the prior treatment with penicillin significantly ($P<0.001$) reduced
607 α -toxin-induced haemolysis. For the penicillin-treated cultures, there were small but significant ($P<0.05$)
608 increases in %H for (a) at 40 and 60mins, and for (b) at 60mins compared to the %H values at 0mins. Data
609 shown are means \pm s.e.m. for at least 3 independent experiments.

610

611 **Figure 6. Comparison between the inhibitory effects of penicillin and linezolid on the extent of α -toxin-**
612 **induced haemolysis.** A single dose of penicillin (250,000U/ml) or linezolid (25 μ g/ml; final concentrations)
613 was added to α -toxin positive strain (DU5946) *S. aureus* cultures after 60mins (corresponding to approximately
614 55% haemolysis) and α -toxin-induced haemolysis determined. There was a significant reduction in levels of
615 α -toxin-induced haemolysis for both drugs at 20, 40, 60 and 80mins compared to the untreated control at the
616 same time points. However the inhibition was significantly greater for the linezolid-treated cultures compared
617 to those following penicillin addition at 40, 60 and 80mins ($P<0.01$). Data shown are means \pm s.e.m. for 3 at
618 least independent experiments.

619

620

621 **Figure 7. Protection of *in situ* chondrocytes by penicillin or linezolid against *S. aureus* infection in a**
622 **bovine model of septic arthritis.** Bovine cartilage was incubated with α -toxin positive strain (DU5946) *S.*
623 *aureus* cultures for 15hrs, which produced approximately 20% chondrocyte death (infected), or in the absence
624 of bacteria (uninfected). Cultures were then continued untreated, or following the addition (at the arrow) of
625 penicillin (1,000U/ml) or linezolid (25 μ g/ml). Data shown are means \pm s.e.m. for 6 independent experiments.
626 Examples of confocal images of chondrocytes in bovine cartilage taken after 23hrs are shown for the following
627 conditions (a) uninfected (<2% cell death), (b) infected penicillin – treated (about 20% cell death), (c) infected
628 linezolid – treated (~21% cell death) and (d) infected untreated (~100% cell death). The scale bar is 100 μ m
629 for all panels.


630

631


632

633


Miller et al., (2018) Fig. 1.



Miller et al., (2018) Fig. 2.




Miller et al., (2018) Fig. 3.




636
637

Miller et al., (2018) Fig. 4.




638
639
640

Miller et al., (2018) Fig. 5.




Miller et al., (2018) Fig. 6.



Miller et al., (2018) Fig. 7.

(a)



(b)

